

Self-similar solutions with fat tails for a coagulation equation with nonlocal drift

Michael Herrmann*

Philippe Laurençot†

Barbara Niethammer‡

January 26, 2009

Abstract

We investigate the existence of self-similar solutions for a coagulation equation with nonlocal drift. In addition to explicitly given exponentially decaying solutions we establish the existence of self-similar profiles with algebraic decay.

1 Introduction

The classical mean-field theory by Lifshitz & Slyozov [5] and Wagner [8] describes the coarsening of droplets in dilute binary mixtures and is based on the assumption that droplets interact only via a common mean-field $u = u(t)$. It results in a nonlocal transport equation for the number density $f = f(t, x) \geq 0$ of droplets of volume $x > 0$ at time $t \geq 0$ and reads

$$\partial_t f(t, x) + \partial_x((a(x)u(t) - b(x))f(t, x)) = 0, \quad \int_0^\infty x f(t, x) dx = m_1 = \text{const}, \quad (1)$$

where the second equation describes the conservation of matter (volume) and determines the mean-field u . The functions a and b are specified by the mechanism of transfer of matter between droplets and a typical example is $a(x) = x^\alpha$ and $b(x) = x^\beta$ with $0 \leq \beta < \alpha \leq 1$ (the original choice in [5] corresponding to $(\alpha, \beta) = (1/3, 0)$).

The large time behaviour of solutions to (1) was conjectured to be given by self-similar solutions already in [5, 8] but it is now known that it actually depends sensitively on the details of the initial data at the end of their support [1, 2, 7]. As a regularisation it was suggested in [5] to add a coagulation term with additive kernel to the evolution equation for f which accounts for the occasional merging of droplets, that is,

$$\begin{aligned} \partial_t f(t, x) &+ \partial_x((a(x)u(t) - b(x))f(t, x)) \\ &= \frac{1}{2} \int_0^x w(x-y, y) f(t, x-y) f(t, y) dy - f(t, x) \int_0^\infty w(x, y) f(t, y) dy, \end{aligned} \quad (2)$$

the mean-field u still being given by the conservation of volume. Well-posedness for this case is proven in [4] and the existence of a fast decaying self-similar solution is established in [3] (for $a(x) = x^{1/3}$, $b(x) = 1$, and $w(x, y) = x + y$); a full characterization of all self-similar solutions seems however difficult, as well as the study of their stability. In fact, these questions are still open for the coagulation equation ($a = b = 0$ in (2)), except for the so-called solvable kernels $w(x, y) = 2$ and $w(x, y) = x + y$ [6]. In particular, besides the existence of exponentially decaying self-similar profiles, nothing is known in general about self-similar solutions with algebraic decay (“fat tails”).

*Oxford Centre for Nonlinear PDE (OxPDE), michael.herrmann@maths.ox.ac.uk

†Institut de Mathématiques de Toulouse, laurenco@math.univ-toulouse.fr

‡Oxford Centre for Nonlinear PDE (OxPDE), niethammer@maths.ox.ac.uk

In order to develop methods to tackle these problems we consider here the following simplified version of (2)

$$\partial_t f(t, x) + \partial_x((xu(t) - 1)f(t, x)) = \int_0^x f(t, x - y)f(t, y) dy - 2f(t, x) \int_0^\infty f(t, y) dy, \quad (3)$$

corresponding to the choice $a(x) = x$, $b(x) = 1$, and $w(x, y) = 2$ for $(x, y) \in (0, \infty)^2$. The function u is again specified by $\int_0^\infty xf(t, x) dx = m_1$, which is equivalent to $u(t) = m_0/m_1$ with $m_0 := \int_0^\infty f(t, x) dx$. Self-similar solutions to (3) are given by $f(t, x) = t^{-2} F(x/t)$ and $u(t) = v/t$, for some $v \in (0, \infty)$. Introducing $z = x/t$ we obtain that (F, v) solves

$$-(z(1-v) + 1)F'(z) = (2-v-2m_0)F(z) + \int_0^z F(z-y)F(y) dy, \quad z \in (0, \infty), \quad (4)$$

where $m_0 = \int_0^\infty F(z) dz$, and v is such that for given $m_1 > 0$ the solution F satisfies

$$\int_0^\infty zF(z) dz = m_1, \quad (5)$$

so (4) implies $v = m_0/m_1$. For the following analysis it is convenient to use m_0 and v as parameters. It is then easily seen that for each $v \in (0, 1)$ there is an exponentially decaying solution

$$F_v(z) = m_0 v e^{-vz} \quad \text{with} \quad m_0 = 1 - v.$$

Besides these self-similar profiles which decay exponentially fast at infinity, we establish the existence of self-similar solutions with algebraic decay provided that the parameter m_0 is sufficiently small.

Theorem 1. *For each $v \in (0, 1)$ there exists $\overline{m_0} \in (0, 1-v)$ such that (4)–(5) has a unique solution F with $\int_0^\infty F(z) dz = m_0$ and $m_1 = m_0/v$ for all $m_0 \in (0, \overline{m_0}]$. This solution F is nonnegative, non-increasing, and satisfies $F(z) \sim cz^{-(2-v)/(1-v)}$ as $z \rightarrow \infty$ for some $c > 0$.*

In fact, we conjecture that for all $m_0 \in (0, 1-v)$ there is a unique solution to (4)–(5) as in Theorem 1 and that there is no solution with $m_0 > 1-v$. We aim to prove this conjecture in a future work by a continuation method starting from the solutions provided by Theorem 1.

2 Proof of the existence result

Within this section we always suppose that $v \in (0, 1)$ is fixed. Furthermore we denote by $M_i(F) := \int_0^\infty y^i F(y) dy$ with $i \in \mathbb{N}$ the i th moment of a nonnegative and integrable function F .

Basic properties of solutions. We start with deriving some elementary properties of solutions of (4)–(5). Integrating (4) we easily establish the following Lemma.

Lemma 2. *Let F be a solution of (4) such that $M_0(F) = m_0$. Then we have $F(0) = m_0(1 - m_0)$, and $m_1 = M_1(F) < \infty$ implies $v = m_0/m_1$.*

This result in particular implies that once we have established the existence of a self-similar solution, then uniqueness follows by uniqueness of the corresponding initial-value problem. Moreover, we also infer from Lemma 2 that positive solutions can only exist for $m_0 \in (0, 1)$, but for technical reasons, and since our results below requires m_0 to be sufficiently small anyway, we assume from now on that $m_0 < v/2$.

Since our existence proof relies on a fixed point argument for F we need appropriate supersolutions for (4). To this end we define $\alpha := (2-v-2m_0)/(1-v) \in (2, \infty)$ and the function $\overline{F}_{m_0}(z) := m_0 (1 + (1-v)z)^{-\alpha}$ for $z \geq 0$, which is the solution to the ordinary differential equation

$$-(z(1-v) + 1)\overline{F}'_{m_0}(z) = (2-v-2m_0)\overline{F}_{m_0}(z), \quad \overline{F}_{m_0}(0) = m_0.$$

This function satisfies $\overline{F}_{m_0}(z) \sim cz^{-\alpha}$ as $z \rightarrow \infty$, and thanks to $m_0 < v/2$ we find $M_1(\overline{F}_{m_0}) < \infty$ as well as $\overline{F}'_{m_0}(z) \leq 0$ for all $z \geq 0$. As a consequence of the maximum principle for ordinary differential equations we readily derive the following comparison result.

Lemma 3. *Any nonnegative solution F to (4) with $F(0) \leq m_0$ satisfies $F(z) \leq \overline{F}_{m_0}(z)$ for all $z \geq 0$.*

From now on we restrict our considerations to *admissible* functions $F \in \mathcal{A}$, where $\mathcal{A}$ is the set of all nonnegative and continuous functions $F : [0, \infty) \rightarrow [0, \infty)$ with finite moments $M_0(F) < \infty$ and $M_1(F) < \infty$.

An auxiliary problem. A key ingredient in our existence proof is to show the existence of solutions to the following auxiliary problem: For a given $G \in \mathcal{A}$ with $M_0(G) = m_0$ we seek $F \in \mathcal{A}$ such that

$$-((1-v)z+1)F'(z) = (2-v-2m_0)F(z) + 2 \int_0^{z/2} F(z-y)G(y) dy, \quad M_0(F) = m_0. \quad (6)$$

Notice that the convolution operator in (6) is related to the integration over the interval $[0, z/2]$. For $F = G$, however, the identity $2 \int_0^{z/2} F(z-y)F(y) dy = \int_0^z F(z-y)F(y) dy$ implies the equivalence of (6) and (4).

Lemma 4. *Consider $G \in \mathcal{A}$ with $M_0(G) = m_0$ and suppose that $F \in \mathcal{A}$ solves (6). Then F is unique, monotonically decreasing, and fulfils $m_0 - 2m_0^2 \leq F(0) \leq m_0$.*

Proof. The uniqueness of solutions follows from the homogeneity of the problem combined with a Gronwall-like argument, and the monotonicity is implied by $F \geq 0$, $G \geq 0$, $v < 1$, and $2-v-2m_0 \geq 0$. To derive the bounds for the initial value we integrate over $z \in (0, \infty)$ to obtain

$$F(0) + (1-v)m_0 = (2-v)m_0 - 2m_0^2 + M_0(F * G) \quad \text{with} \quad (F * G)(z) := 2 \int_0^{z/2} F(z-y)G(y) dy,$$

and the estimates

$$0 \leq M_0(F * G) = 2 \int_0^\infty \int_0^{z/2} F(z-y)G(y) dy dz = 2 \int_0^\infty \int_0^z G(y)F(z) dy dz \leq 2m_0^2$$

completes the proof. $\square$

For the subsequent considerations it is convenient to introduce the function $\tau : [0, \infty) \rightarrow \mathbb{R}$ defined by

$$\tau(z) := -z \frac{F'(z)}{F(z)} \quad \text{as long as} \quad F(z) > 0, \quad (7)$$

so that

$$F(z) = F(0) \exp \left(- \int_0^z \frac{\tau(s)}{s} ds \right). \quad (8)$$

Notice in particular that $\tau(z) \rightarrow \tau_\infty \neq 0$ as $z \rightarrow \infty$ implies $F(z) \sim cz^{-\tau_\infty}$ as $z \rightarrow \infty$. Rewriting (6) in terms of τ yields

$$\frac{(1-v)z+1}{z} \tau(z) = 2-v-2m_0 + h[G, \tau](z), \quad h[G, \tau](z) := 2 \int_0^{z/2} G(y) \exp \left(\int_{z-y}^z \frac{\tau(s)}{s} ds \right) dy, \quad (9)$$

and this is equivalent to the fixed point equation

$$\tau = H[G, \tau], \quad H[G, \tau](z) := \frac{z}{(1-v)z+1} (2-v-2m_0 + h[G, \tau](z)). \quad (10)$$

In the following Lemma we summarize some useful properties of the operator H .

Lemma 5. For each $G \in \mathcal{A}$ the operator $\tau \mapsto H[G, \tau]$ is well defined on the set $C([0, \infty))$ and has the following properties:

1. $0 \leq H[G, \tau]$ for all τ ,
2. $\tau_1 \leq \tau_2$ implies $H[G, \tau_1] \leq H[G, \tau_2]$,
3. $\tau(z) \rightarrow \tau_\infty$ as $z \rightarrow \infty$ implies $H[G, \tau](z) \rightarrow (2-v)/(1-v)$ as $z \rightarrow \infty$.

Proof. The first two claims are direct consequences of $G \geq 0$ combined with $v < 1$ and $m_0 \leq v/2 < 1/2$, and the definitions (9) and (10). Now suppose that $\tau(z) \rightarrow \tau_\infty$, and write

$$h[G, \tau](z) = 2 \int_0^{z/2} G(y) \exp \left(\int_{z-y}^z \frac{\tau(s) - \tau_\infty}{s} ds \right) \left(\frac{z}{z-y} \right)^{\tau_\infty} dy.$$

Due to $y \leq z/2$ the convergence assumption implies

$$\left| \exp \left(\int_{z-y}^z \frac{\tau(s) - \tau_\infty}{s} ds \right) - 1 \right| \leq \exp \left(\ln 2 \sup_{s \geq z/2} |\tau(s) - \tau_\infty| \right) - 1 \xrightarrow{z \rightarrow \infty} 0,$$

and from $M_0(G) = m_0 < \infty$, the inequality $z/(z-y) \leq 2$ for $y \in (0, z/2)$, and the Lebesgue Dominated Convergence Theorem we infer that

$$\lim_{z \rightarrow \infty} 2 \int_0^{z/2} G(y) \left(\frac{z}{z-y} \right)^{\tau_\infty} dy = 2 \int_0^\infty G(y) dy = 2m_0.$$

Hence, we have $h[G, \tau](z) \rightarrow 2m_0$ as $z \rightarrow \infty$, and the proof is complete. $\square$

Solutions to the auxiliary problem. In order to set up an iteration scheme for τ we prove that for all sufficiently small m_0 there exists a supersolution to the fixed point equation (10). Notice that the constants in the next Lemma depend on v .

Lemma 6. There exist two constants $\overline{m}_0 \in (0, v/2)$ and $\tau_\star \in \mathbb{R}$ such that $H[G, \tau_\star] \leq \tau_\star$ holds for all $G \in \mathcal{A}$ with $m_0 = M_0(G) \leq \overline{m}_0$.

Proof. The definition (9) gives

$$h[G, \tau](z) \leq 2 \int_0^{z/2} G(y) \left(\frac{z}{z-y} \right)^{\|\tau\|_\infty} dy \leq 2m_0 2^{\|\tau\|_\infty}$$

and hence $H[G, \tau] \leq a_{m_0} + b_{m_0} 2^{\|\tau\|_\infty}$ with $a_{m_0} := (2-v-2m_0)/(1-v)$ and $b_{m_0} := 2m_0/(1-v)$. Since $b_{m_0} \rightarrow 0$ as $m_0 \rightarrow 0$ we choose $\overline{m}_0 > 0$ and $\sigma_\star > 0$ such that $b_{\overline{m}_0} 2^{a_0} \leq \sigma_\star 2^{-\sigma_\star}$, and set $\tau_\star = a_0 + \sigma_\star$. Then, for all $m_0 \in (0, \overline{m}_0]$ we find $a_{m_0} + b_{m_0} 2^{\tau_\star} \leq a_0 + b_{\overline{m}_0} 2^{a_0} 2^{\sigma_\star} \leq a_0 + \sigma_\star = \tau_\star$, and this implies the claimed result. $\square$

Now we are able to prove the existence of solutions to the auxiliary problem.

Corollary 7. For each $G \in \mathcal{A}$ with $m_0 = M_0(G) \in (0, \overline{m}_0]$ there exists a unique solution $F \in \mathcal{A}$ to (6) such that the function τ given by (7) satisfies $0 \leq \tau(z) \leq \tau_\star$ for all $z \geq 0$ and $\tau(z) \rightarrow (2-v)/(1-v)$ as $z \rightarrow \infty$. In addition, $F \leq \overline{F}_{m_0}$.

Proof. We define a sequence of continuous functions $(\tau_i)_i$ by $\tau_0 = \tau_\star$ and $\tau_{i+1} = H[G, \tau_i]$, which satisfies $0 \leq \tau_{i+1} \leq \tau_i \leq \tau_\star$ thanks to Lemma 5 and Lemma 6. Consequently, this sequence converges pointwisely to a function τ , which solves the fixed point equation (10), and is hence continuous. The function F is then determined by (8) and the condition $M_0(F) = m_0$, the latter being meaningful as the behaviour of τ as $z \rightarrow \infty$ guarantees the integrability of F . Moreover, F satisfies the first equation in (6) by construction and $F \leq \overline{F}_{m_0}$ due to Lemma 4, and this implies $M_1(F) < \infty$. $\square$

Fixed point argument for F . We finish the proof of Theorem 1 by applying Schauder's Fixed Point Theorem. For $m_0 \in (0, \overline{m}_0]$ let $\mathcal{M}_{m_0}$ denote the set of all functions $G \in \mathcal{A}$ that satisfy $M_0(G) = m_0$ and $G \leq \overline{F}_{m_0}$. In view of Lemma 4 and Corollary 7, we may define an operator $\mathcal{S}_{m_0} : \mathcal{M}_{m_0} \rightarrow \mathcal{M}_{m_0}$ as follows: For given $G \in \mathcal{M}_{m_0}$ the function $\mathcal{S}_{m_0}[G]$ is the solution to the auxiliary problem (6) with datum G .

Lemma 8. *For each $m_0 \in (0, \overline{m}_0]$ the operator $\mathcal{S}_{m_0}$ has a unique fixed point in $\mathcal{M}_{m_0}$, which satisfies both (4) and (5) with $m_1 = m_0/v$.*

Proof. From (6) and Lemma 4 we infer that each $F \in \mathcal{S}[\mathcal{M}_{m_0}]$ is differentiable with

$$\|F'\|_\infty \leq (2 - v + 2m_0)\|F\|_\infty \leq (2 - v + 2m_0)m_0.$$

Therefore, $\mathcal{S}_{m_0}[\mathcal{M}_{m_0}]$ is precompact in $C([0, \infty))$, and due to the uniform supersolution $\overline{F}_{m_0}$ we readily verify that the integral constraint $M_0(G) = m_0$ is preserved under strong convergence in $\mathcal{M}_{m_0}$. Schauder's theorem now implies the existence of a fixed point $F = \mathcal{S}_{m_0}[F] \in \mathcal{M}_{m_0}$, which satisfies (4) and $M_0(F) = m_0$ by construction. Moreover, $F \leq \overline{F}_{m_0}$ implies $M_1(F) < \infty$, so Lemma 2 guarantees (5). Finally, the fixed point is unique as discussed in the remark to Lemma 2. $\square$

References

- [1] J. Carr. Stability of self-similar solutions in a simplified LSW model. *Phys. D*, 222(1-2):73–79, 2006.
- [2] J. Carr and O. Penrose. Asymptotic behaviour in a simplified Lifshitz–Slyozov equation. *Phys. D*, 124:166–176, 1998.
- [3] M. Herrmann, B. Niethammer, and J.J.L. Velázquez. Self-similar solutions for the LSW model with encounters. 2007. preprint.
- [4] Ph. Laurençot. The Lifshitz–Slyozov equation with encounters. *Math. Models Methods Appl. Sci.*, 11(4):731–748, 2001.
- [5] I. M. Lifshitz and V. V. Slyozov. The kinetics of precipitation from supersaturated solid solutions. *J. Phys. Chem. Solids*, 19:35–50, 1961.
- [6] G. Menon and R. L. Pego. Approach to self-similarity in Smoluchowski's coagulation equations. *Comm. Pure Appl. Math.*, 57(9):1197–1232, 2004.
- [7] B. Niethammer and R. L. Pego. Non-self-similar behavior in the LSW theory of Ostwald ripening. *J. Statist. Phys.*, 95(5/6):867–902, 1999.
- [8] C. Wagner. Theorie der Alterung von Niederschlägen durch Umlösen (Ostwald-Reifung). *Z. Elektrochem.*, 65:581–591, 1961.